

About the Authors



Akhil Rajendra, MD, DM

Dr. Akhil Rajendra is a Clinical Research fellow in the Leukemia and Myeloproliferative neoplasms program at the Princess Margaret Cancer Centre, University of Toronto. Prior to coming to Toronto, he obtained his M.D. in internal medicine at Christian Medical College, Vellore, India. He completed his subspecialty training in medical oncology and haemato-oncology at Tata Memorial Hospital, Mumbai, India. He has a special interest in acute leukemias and myeloid disorders.

Affiliations: Clinical Research Fellow, Leukemia and Myeloproliferative Neoplasm, Division of Medical Oncology & Hematology, Princess Margaret Cancer Centre, University Health Network, University of Toronto



Dawn Maze, MD, FRCPC, MSc

Dr. Dawn Maze is a clinician investigator at the Princess Margaret Cancer Centre and an Assistant Professor in the Department of Medicine at the University of Toronto. She is a member of the Leukemia Site Group and the Myeloproliferative Neoplasms (MPN) Program and has a clinical research interest in myeloid malignancies. Her primary research focus is in MPN, particularly in risk stratification and management approaches early in the disease course and in the adolescent/young adult population. She is particularly interested in optimizing pregnancy outcomes in this population. She is also actively involved in clinical trials and projects aiming to refine risk stratification in acute myeloid leukemia. Dr. Maze is an active member of the Canadian MPN Group. She is the Medical Director for the Malignant Hematology Day Unit at Princess Margaret and the Program Director for the Leukemia Fellowship Program.

Affiliations: Division of medical oncology and hematology, Princess Margaret Cancer Centre, University of Toronto, Toronto, ON, Canada

Treatment of Philadelphia Chromosome-negative Myeloproliferative Neoplasms in 2024: A Concise Review

Akhil Rajendra, MD, DM
Dawn Maze, MD, FRCPC, MSc

Introduction

In 1951, William Dameshek coined the term myeloproliferative disorders (MPDs) for diseases characterized by abnormal proliferation of one or more terminally differentiated myeloid cell lines in the peripheral blood.^{1,2} In 2008, the World Health Organization (WHO) renamed these disorders as myeloproliferative neoplasms (MPNs) in recognition of their clonal nature. There are currently two classification system for MPNs: WHO and International Consensus Classification (ICC), 2022.^{3,4} This review will focus on the Philadelphia chromosome-negative MPNs, which include polycythemia vera (PV), essential thrombocythemia (ET), and primary myelofibrosis (PMF).

Genomic changes in MPNs

MPNs result from the constitutive activation of the Janus kinase/signal transducer and activator of transcription (JAK/STAT) signalling pathway. The JAK2 p.V617F mutation, first described in 2005, is detectable in >95% of patients with PV and 50-60% of patients with ET or PMF. In-frame insertions or deletions in exon 12 of the JAK2 gene are found in the remaining patients with PV but not in those with ET.⁵⁻⁷ Mutations in the thrombopoietin receptor gene *MPL* were identified in 2006 and are present in 3-5% of ET and 5-10% of PMF, but not in PV cases.⁸ Mutations in the calreticulin (*CALR*) gene were identified in 2013 and are found in 20-25% of ET and 25-30% of PMF but not in PV.^{9,10} The *CALR* gene encodes for the endoplasmic reticulum chaperone protein (*CALR*). Mutant *CALR* interacts with the *MPL* protein, which is trafficked to the cell surface thereby activating

the JAK-STAT signalling pathway.¹¹ Mutations in the *CALR* gene consist of insertions or deletions in exon 9 resulting in a positively charged amino acid sequence in the C-terminus. The mutations can be type 1, characterised by a 52-bp deletion that eliminates all the negatively charged amino acids in the C-terminus, or type 2, characterised by 5-bp insertion that eliminates half the negatively charged amino acids from the C-terminus. Type 1 and type 2 mutations constitute 80% of the *CALR* mutations.

In addition to the above three driver mutations, other somatic myeloid mutations are also found in MPNs. Common somatic mutations involve genes regulating DNA methylation (*TET2*, *DNMT3A*, and *IDH1/IDH2*), histone modification (*ASXL1* and *EZH2*), RNA splicing (*SF3B1*, *U2AF1*, *ZRSR2*, and *SRSF2*), and the RAS pathway (*NRAS* and *KRAS*). These mutations are common in PMF and the blast phase of PV and ET. While these mutations do not cause MPN, they may modify the disease phenotype. Mutations in *ASXL1*, *EZH2*, *SRSF2*, *U2AF1*, and *IDH1/2* are denoted as resulting in the “high molecular risk” phenotype.¹²

Management of PV

PV is a clonal hematopoietic stem cell neoplasm characterized by panmyelosis, disease-related symptoms, increased risk for thrombosis, and risk of transformation to post-PV myelofibrosis (MF) or acute leukemia. Goals of treatment for PV include prevention of thrombosis, reducing symptom burden, and prevention of disease progression.

PV-related thrombosis is multifactorial and related to hyperviscosity, increased red cell mass, and increased thrombin generation by platelets.¹³

JAK2 positivity contributes to thrombosis risk in MPN¹⁴, as does increased allele burden.¹⁵ Once-daily aspirin (81 mg/day; acetylsalicylic acid [ASA]) is recommended for all patients with PV without contraindications.¹⁶ In addition, phlebotomies are performed to achieve a target hematocrit level of <45%.^{17,18}

Beyond phlebotomy and aspirin, cytoreductive treatment is indicated for individuals with high-risk disease.¹⁹ Traditionally, patients who are over 60 years of age and/or have a history of thrombosis are considered to have high-risk disease, while those without these factors are considered low risk.¹⁹ In certain scenarios cytoreductive therapy may be considered even in patients with low-risk disease (**Figure 1**):

1. Frequent phlebotomies with suboptimal hematocrit control or poor tolerability
2. Symptoms of PV (microvascular, pruritis) not controlled with ASA or phlebotomies
3. Phlebotomies leading to symptomatic iron deficiency anemia
4. Extreme thrombocytosis leading to acquired von Willebrand syndrome

Cytoreductive therapy

Over the years, hydroxyurea (HU) has been the standard cytoreductive agent in PV. HU is usually started at a dose of 500 mg once or twice daily, and titrated based on response. Another option, interferon alfa (IFN α), has long been shown to have cytoreductive and disease-modifying potential. However, its toxicity and need for frequent parenteral administration has been a deterrent to its usage. This has changed with the availability of pegylated forms of IFN α . The only formulation currently available in Canada is peginterferon alfa-2a (Pegasys). Another formulation is ropeginterferon alfa-2b (rIFN), which is a monopegylated form of IFN α . This formulation is characterised by an extended elimination half-life, resulting in less frequent dosing, better tolerability, and improved compliance.²⁰ This formulation is FDA-approved.

Phase 3 trials have established the role of IFN α in high-risk PV. The MPD-RC-112 trial, in which randomized patients with high-risk ET/PV received Pegasys or HU²¹, and the PROUD-PV and CONTINUATION PV studies randomized patients

with high-risk PV to receive rIFN or HU.^{22,34} IFN α was non-inferior to HU in terms of complete hematological response (CHR) at 12 months in both these trials.²¹⁻²³ In the CONTINUATION-PV study, CHR was higher for the rIFN group in long-term follow-up.²³ JAK2 allele burden decreased consistently over time with both IFN α drugs, which was associated with improved event-free survival (EFS).²⁴ The starting dose for Pegasys is 45 mcg subcutaneously weekly. Doses are titrated with 45 mcg monthly increments to a maximum of 180 mcg.²¹ rIFN is administered subcutaneously every 2 weeks at a starting dose of 100 or 50 mcg (for HU-exposed patients). Dosing increments of 50 mcg are made every 2 weeks up to a maximum of 500 mcg.^{22,25}

Treatment of patients with HU-intolerant, resistant disease

A significant number of patients are intolerant to HU due to hematologic or non-hematologic toxicity or their disease is resistant to this therapy due to a lack of effective cytoreduction. HU intolerance or resistance has been defined by the European LeukemiaNet (ELN; **Table 1**)^{26,27}:

1. Need for phlebotomy to maintain hematocrit levels <45% after 3 months of at least 2 g/day of hydroxyurea **OR**
2. Uncontrolled myeloproliferation (i.e. platelet count >400 $\times 10^9/L$ and white blood cell count >10 $\times 10^9/L$) after 3 months of at least 2 g/day of hydroxyurea **OR**
3. Failure to reduce massive splenomegaly by more than 50% as measured by palpation or failure to completely relieve symptoms related to splenomegaly, after 3 months of at least 2 g/day of hydroxyurea **OR**
4. Absolute neutrophil count <1.0 $\times 10^9/L$, or platelet count <100 $\times 10^9/L$, or hemoglobin <100 g/L at the lowest dose of hydroxyurea required to achieve complete or partial clinical hematological response **OR**
5. Presence of leg ulcers or other hydroxyurea-related non-hematological toxicities like mucocutaneous manifestations, gastrointestinal symptoms, pneumonitis, or fever at any dose of hydroxyurea

Table 1. Definition of clinical resistance and intolerance to hydroxyurea in polycythemia vera and myelofibrosis; adapted from Barosi, G, et al., 2007 and Barosi, G, et al., 2010.




Figure 1. Approach to management of polycythemia vera; adapted from Tefferi and Barbui, 2008.¹⁹

In the MPD-RC-111 trial, a single-arm Phase 2 study, patients with HU-resistant or -intolerant disease were treated with Pegasys, which resulted in a 12-month overall response rate (ORR) of 60% and spleen normalisation in 32.7% of cases.²⁸ Ruxolitinib, an oral JAK inhibitor, was also assessed in this population in three randomized trials: the RESPONSE trial (with splenomegaly)²⁹, the RESPONSE-2 trial (without splenomegaly)³⁰, and the MAJIC-PV study (Phase 2). The comparator arm in these trials was the best available therapy (BAT). All three trials showed that ruxolitinib was better at achieving hematocrit control and spleen volume reduction compared to BAT. The MAJIC-PV trial also showed better EFS with ruxolitinib.³¹ However, IFN α -based therapy constituted only 11.6%, 13%, and 15% of BATs.²⁹⁻³¹ Thus, whether ruxolitinib or pegIFN α is the best agent for those with HU-resistant/intolerant disease remains unknown. Future trials must focus on the appropriate sequencing of these agents for this group of patients.

Novel approaches

IFN α in low-risk PV

The role of rIFN in low-risk PV was studied in the LOW PV study which was a Phase 2 randomized trial comparing rIFN with phlebotomy. The group receiving rIFN had better hematologic response,^{32,33} (rIFN was dosed 100 mcg every 2 weeks with no escalation).

Hepcidin-mimetic (rusfertide) in PV

Hepcidin binds to ferroportin, blocking the export of intracellular iron to the blood leading to reduced serum iron levels and decreased erythropoiesis.³⁴ In the Phase 2 REVIVE trial involving patients with phlebotomy-dependent PV, rusfertide was associated with a significant decline in phlebotomies and better hematological response.³⁵ The ongoing Phase 3 VERIFY trial is evaluating its efficacy and safety in PV.³⁶

In summary, patients with low-risk PV are managed with aspirin and phlebotomy to achieve hematocrit levels of <45%. Cytoreductive therapy is indicated in patients with high-risk PV. In certain scenarios in low-risk PV, cytoreductive therapy can be instituted. Both the National Comprehensive Cancer Network (NCCN) and ELN recommend either HU or pegIFN α /rIFN as first line cytoreductive therapies. pegIFN α or rIFN are favoured in younger patients (<60 years) and women of child-bearing age.³⁷ In the

HU-resistant/intolerant population, both pegIFN and ruxolitinib can be used.

Management of ET

ET is characterised by predominantly thrombocytosis, occurrence of thrombosis, and microcirculatory symptoms, and occasionally disease transformation to fibrosis or leukemia.

Risk-stratified treatment

Similar to PV, treatment in ET is focused on thrombosis prevention. Traditional risk factors include age over 60 years and history of thrombosis.³⁸ More recently, the international prognostic score for ET (IPSET), has refined risk stratification in ET by incorporating JAK2 mutation status. In its latest iteration the revised IPSET thrombosis score categorises patients into four risk groups (**Table 2**).^{39,40}

Despite the lack of randomized evidence, low-dose aspirin is used for thrombosis prevention in ET. Recommendations are based on non-randomized studies^{41,42} and by extrapolation from studies in PV.¹⁶ In the absence of contraindications, low-dose aspirin is a reasonable choice in patients with low, intermediate, and high-risk disease and in those with very low-risk disease with microvascular symptoms. In a recent study of low-risk patients with mutated *CALR*, no benefit was observed for the use of low-dose aspirin, while it was associated with increased risk of bleeding.⁴¹ In patients with extreme thrombocytosis ($>1000 \times 10^9/L$), aspirin should be used with caution due to the risk of bleeding and acquired von Willebrand factor deficiency (**Figure 2**).

Cytoreductive therapy

The first line cytoreductive therapy of choice for ET is HU. Similar to PV, pegylated IFN can be used in ET. The MPD-RC-112 trial compared Pegasys with HU in high-risk ET. The percentage of patients with complete remissions (CR) at 12 months were 44% and 45% with Pegasys and HU, respectively.²¹ Anagrelide, an oral imidazoquinoline, when compared with HU in the first line, resulted in higher rates of thrombosis (arterial and venous), hemorrhage, and transformation to myelofibrosis than HU.⁴³

For the HU intolerant/resistant population, the MPD-RC-111 trial showed that Pegasys produces reasonable responses (ORR of 69% at 12 months).²⁸ On the other hand, in the

Risk	Attributes	Management
Very low	Age ≤60 years, JAK2 wild type, no prior thrombosis	Observation Low-dose aspirin (in the presence of cardiovascular risk factors)
Low	Age ≤60 years, JAK2 V617F mutated, no prior thrombosis	Low-dose aspirin
Intermediate	Age >60 years, JAK2 V617F wild type, no prior thrombosis	Low-dose aspirin +/- cytoreductive therapy
High	Age >60 years and JAK2 V617F mutated OR Prior thrombosis regardless of other factors	Low-dose aspirin + cytoreductive therapy

Table 2. Revised international prognostic score for ET; adapted from Barbui et al., 2015.⁴⁶

MAJIC-ET trial, when ruxolitinib was compared to BAT in this population, the ORR, and rates of thrombosis, hemorrhage, and transformation were similar. The BAT used were IFNα, anagrelide, busulfan, and HU.⁴⁴

Thus, in patients with high-risk ET, the first line cytoreductive therapy of choice is HU. Pegylated IFN should be considered in younger patients and individuals of child-bearing age. Either of these agents (HU or IFNα) can be used in the second line if not previously used and anagrelide is an alternative option. Ruxolitinib has activity in ET and may be considered in certain circumstances. Results of the SURPASS-ET trial, comparing ruxolitinib with anagrelide in HU-intolerant/resistant ET are pending.⁴⁵

Treatment of PMF and post-PV/ET MF

Primary myelofibrosis (PMF) is characterised by progressive cytopenia, marrow fibrosis, cytokine-driven inflammatory symptoms, and extramedullary hematopoiesis. A disease phenotype similar to PMF is observed in advanced phases of PV and ET and is defined as post-PV-MF and post-ET-MF, respectively. Aberration in the JAK/STAT signaling pathways is crucial to the pathogenesis of MF, which in 90% of patients is driven by mutually exclusive mutations in JAK2, CALR, or MPL genes.⁴⁷ Somatic mutations in the myeloid genes (mentioned under genomic changes) additionally influence MF biology.⁴⁸

Risk stratification

Management of MF begins with risk stratification. Earlier risk models include the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), and DIPSS-plus.⁴⁹⁻⁵¹ Better genomic understanding has led to the incorporation of genetic mutations into the risk stratification. Mutations in ASXL1, SRSF2, IDH1/2, and EZH2 confer poorer prognosis.⁵¹ Mutational data has been integrated into the Mutation-enhanced (M) IPSS70, MIPSS70-plus, and MIPSS70+ version 2.0 risk stratification models.^{53,54} Mutations in the TP53 gene are not included in these risk systems. Seminal work by Grinfeld et al. showed that TP53-mutated MF has a high risk of leukemic transformation and very poor median overall survival (OS) of 2.4 years.⁴⁸ These risk models have been validated in primary myelofibrosis but not in secondary myelofibrosis. In clinical practice, these models are frequently used in secondary MF. The Myelofibrosis Secondary to PV and ET – Prognostic Model (MYSEC-PM) is a prognostic model developed specifically for secondary MF.⁵⁵




Figure 2. Approach to management of essential thrombocythemia; adapted from Barbui et al., 2015.⁴⁶

Treatment of MF

Patients with DIPSS scores intermediate 2 or higher, MIPSS70 or MIPSS70-plus version 2.0 high risk, MYSEC-PM intermediate 2 or higher, and *TP53* mutations have a predicted median overall survival of <5 years and should be considered for allogeneic stem cell transplantation (**Figure 3**).⁵⁶ Peri-transplant management is directed at symptoms and splenomegaly and a bridging JAK inhibitor (JAKi) can be considered. Timing of the transplant in the JAKi era is controversial and is covered in other publications.⁵⁷⁻⁵⁹ For patients who are ineligible for transplant, do not have a suitable donor, or prefer non-transplant therapy, JAKi have been the mainstay of therapy for symptomatic management. Patients who are not high risk per the above models can be monitored if asymptomatic,

receive symptom-directed management, or referred to clinical trials as appropriate.

Choice of JAKi

There are currently four FDA-approved JAKi for myelofibrosis: ruxolitinib, fedratinib, pacritinib, and momelotinib, the first two of which are Health Canada approved. Ruxolitinib, a non-selective JAK1/JAK2 inhibitor, approved in the US in 2011 and in Canada in 2012, has the largest body of evidence. In the COMFORT-I and COMFORT-II trials comparing ruxolitinib to placebo and BAT, respectively, ruxolitinib resulted in a spleen volume reduction of 35% (SVR35) at 24 weeks (SVR35@24) in 41.9% and 32% of patients, respectively.^{60,61} Anemia and thrombocytopenia are important side effects




Figure 3. Management algorithm for transplant-eligible patients with MF in the chronic phase; *used with permission from Davidson and Gupta, 2021.*⁵⁸

of ruxolitinib, which lead to dose reductions or treatment interruptions. At 3 years, 50% of patients had discontinued ruxolitinib, and this rate increased to 75% at 5 years.⁶²

Fedratinib is a JAK2-FLT3-BRD4 inhibitor that has been studied in both ruxolitinib-naïve and -exposed patients in the JAKARTA and JAKARTA-2 trials. To be included in these trials, platelet levels had to be $\geq 50 \times 10^9/L$. Fedratinib resulted in a SVR35@24 of 36% and 55%, respectively, with good symptom burden reduction.⁶³⁻⁶⁶ Even though fedratinib is effective in the first line setting, ruxolitinib is most often used in clinical practice. The Health Canada approval for fedratinib is for patients with MF with disease-related symptoms or splenomegaly, including those who have been previously exposed to ruxolitinib.⁶⁷

Momelotinib is a JAK1/JAK2 inhibitor that has additional inhibitory effects against activin A receptor type 1 (ACVR1). ACVR1 is involved in

SMAD2/3 signalling, which upregulates hepcidin production. Momelotinib has been found to have significant anemia benefits. In the SIMPLIFY-1 trial, momelotinib was found to be non-inferior to ruxolitinib in terms of the SVR35@24, but not for symptom score reduction.⁶⁸ In addition, this trial showed that red blood cell (RBC) transfusion independence and conversion to transfusion independence was better with momelotinib.⁶⁹ Momelotinib is an exciting option for the treatment of symptomatic MF with anemia. Approval in Canada is anticipated in the near future.

The fourth JAKi is pacritinib, which was studied in the PERSIST-1 and PERSIST-2 trials that included patients with platelet counts $< 50 \times 10^9/L$ (both JAKi-naïve and JAKi-exposed). Pacritinib achieved SVR35@24 of 23.1% and symptom control in 25% of patients.⁷⁰

Combination therapy

A number of novel agents have been combined with JAKi therapy in clinical trials. In the MANIFEST-2 trial, patients with treatment-naïve MF with an enlarged spleen (DIPSS intermediate-1 or higher) were randomized to receive ruxolitinib + pelabresib (BET inhibitor) or ruxolitinib + placebo.⁷¹ In the TRANSFORM-1 trial, the combination of ruxolitinib + navitoclax (BCL-2 inhibitor) was compared with ruxolitinib + placebo.⁷² Both combinations resulted in a doubling of the SVR35@24 in comparison to ruxolitinib + placebo. However, neither combinations significantly reduced the symptom burden in comparison to ruxolitinib + placebo. Therefore, the place of these combinations for treatment remains unclear and longer follow-up studies are awaited. These two trials also highlight the need for better endpoints to evaluate therapies in MF. In addition, the combination of ruxolitinib + pelabresib showed improvement of bone marrow fibrosis.⁷³ This could be evidence of disease modification with the BET inhibitor. Other therapies with disease-modifying potential are required.

Agents addressing anemia

Transfusion dependence is a major symptom in MF. Transfusion dependence is associated with poorer overall survival in patients with MF.^{74,75} Apart from momelotinib and pacritinib, which positively affect anemia due to *ACVR1* inhibition, there are other adjunctive therapies that have been used in patients with MF and anemia. RBC transfusion is the most commonly used strategy in clinical practice. Erythropoietin (EPO)-stimulating agents can be used in patients with EPO levels <500 U/L with an expected response ranging from 40–60%.^{76–78} Androgens (danazol), steroids, immunomodulatory agents (lenalidomide, thalidomide), and splenectomy are other strategies that have been used.⁷⁹ Recently, the Phase 2 open label ACE-536-MF-001 trial tested luspatercept in patients with MF. Luspatercept resulted in improvement of the primary endpoint (anemia response) in transfusion-dependent (9.5%) and non-transfusion-dependent (13.6%) patients and in patients who were on ruxolitinib (26.3% and 14.3%, respectively).⁸⁰

In summary, management of MF begins with risk stratification. Patients with high-risk disease should be offered a transplant. JAKi can be used in peri-transplant symptom

management. In patients who are ineligible for transplant or decline transplant, management is symptom-directed using JAKi. Ruxolitinib is the JAKi with the most clinical experience. Newer JAKi, such as momelotinib and pacritinib, have a role in the setting of co-existing cytopenia. Trials are assessing agents that modify the disease biology and also address anemia.

Conclusions and future directions

The past decade has seen major shifts in the diagnosis, prognostication, and management of MPN. The focus of treatment for PV and ET is thrombosis prevention and monitoring for disease progression. New data support the use of IFN α therapy for cytoreduction, especially in PV, and also appears to result in sustained decline in JAK2 allele burden in a proportion of patients. Management of MF begins with risk assessment. Patients with high-risk disease should be considered for transplant. Symptom management of MF has seen the availability of several JAK inhibitors which may help address the co-existing cytopenia in MF. With the availability of many agents, sequencing of therapies will become increasingly important in the future. Several agents are focused on addressing anemia in MF, which continues to be an area of unmet need. Patients should be offered clinical trial participation whenever possible.

Disclaimer: At the time of publishing this review, there is a global shortage in the supply of Pegasys, which is expected to last until the second half of 2025.

Correspondence

Dawn Maze, MD, FRCPC, MSc
Email: dawn.maze@uhn.ca

Financial Disclosures

A.R.: None declared.

D.M.: None declared.

References

1. Tefferi A. The history of myeloproliferative disorders: before and after Dameshek. *Leukemia*. 2008 Jan;22(1):3–13.
2. DAMESHEK W. Editorial: Some Speculations on the Myeloproliferative Syndromes. *Blood*. 1951 Apr 1;6(4):372–5.

3. Khoury JD, Solary E, Ablu O, Akkari Y, Alaggio R, Apperley JF, et al. The 5th edition of the World Health Organization Classification of Haematolymphoid Tumours: Myeloid and Histiocytic/Dendritic Neoplasms. *Leukemia*. 2022 Jul;36(7):1703–19.
4. Arber DA, Orazi A, Hasserjian RP, Borowitz MJ, Calvo KR, Kvasnicka HM, et al. International Consensus Classification of Myeloid Neoplasms and Acute Leukemias: integrating morphologic, clinical, and genomic data. *Blood*. 2022 Sep 15;140(11):1200–28.
5. Baxter EJ, Scott LM, Campbell PJ, East C, Fourouclas N, Swanton S, et al. Acquired mutation of the tyrosine kinase JAK2 in human myeloproliferative disorders. *Lancet Lond Engl*. 2005 Mar 19;365(9464):1054–61.
6. Mejía-Ochoa M, Acevedo Toro PA, Cardona-Arias JA. Systematization of analytical studies of polycythemia vera, essential thrombocythemia and primary myelofibrosis, and a meta-analysis of the frequency of JAK2, CALR and MPL mutations: 2000–2018. *BMC Cancer*. 2019 Jun 17;19(1):590.
7. Scott LM, Tong W, Levine RL, Scott MA, Beer PA, Stratton MR, et al. JAK2 exon 12 mutations in polycythemia vera and idiopathic erythrocytosis. *N Engl J Med*. 2007 Feb 1;356(5):459–68.
8. Staerk J, Lacout C, Sato T, Smith SO, Vainchenker W, Constantinescu SN. An amphipathic motif at the transmembrane-cytoplasmic junction prevents autonomous activation of the thrombopoietin receptor. *Blood*. 2006 Mar 1;107(5):1864–71.
9. Klampfl T, Gisslinger H, Harutyunyan AS, Nivarthi H, Rumi E, Milosevic JD, et al. Somatic Mutations of Calreticulin in Myeloproliferative Neoplasms. *N Engl J Med*. 2013 Dec 19;369(25):2379–90.
10. Nangalia J, Massie CE, Baxter EJ, Nice FL, Gundem G, Wedge DC, et al. Somatic CALR Mutations in Myeloproliferative Neoplasms with Nonmutated JAK2. *N Engl J Med*. 2013 Dec 19;369(25):2391–405.
11. How J, Hobbs GS, Mullally A. Mutant calreticulin in myeloproliferative neoplasms. *Blood*. 2019 Dec 19;134(25):2242–8.
12. Lundberg P, Karow A, Nienhold R, Looser R, Hao-Shen H, Nissen I, et al. Clonal evolution and clinical correlates of somatic mutations in myeloproliferative neoplasms. *Blood*. 2014 Apr 3;123(14):2220–8.
13. Barbui T, Finazzi G, Falanga A. Myeloproliferative neoplasms and thrombosis. *Blood*. 2013 Sep 26;122(13):2176–84.
14. Campbell PJ, Scott LM, Buck G, Wheatley K, East CL, Marsden JT, et al. Definition of subtypes of essential thrombocythemia and relation to polycythemia vera based on JAK2 V617F mutation status: a prospective study. *The Lancet*. 2005 Dec 3;366(9501):1945–53.
15. Borowczyk M, Wojtaszewska M, Lewandowski K, Gil L, Lewandowska M, Lehmann-Kopydłowska A, et al. The JAK2 V617F mutational status and allele burden may be related with the risk of venous thromboembolic events in patients with Philadelphia-negative myeloproliferative neoplasms. *Thromb Res*. 2015 Feb;135(2):272–80.
16. Landolfi R, Marchioli R, Kutti J, Gisslinger H, Tognoni G, Patrono C, et al. Efficacy and Safety of Low-Dose Aspirin in Polycythemia Vera. *N Engl J Med*. 2004 Jan 8;350(2):114–24.
17. Barbui T, Passamonti F, Accorsi P, Pane F, Vannucchi AM, Velati C, et al. Evidence- and consensus-based recommendations for phlebotomy in polycythemia vera. *Leukemia*. 2018 Sep;32(9):2077–81.
18. Marchioli R, Finazzi G, Specchia G, Cacciola R, Cavazzina R, Cilloni D, et al. Cardiovascular Events and Intensity of Treatment in Polycythemia Vera. *N Engl J Med*. 2013 Jan 3;368(1):22–33.
19. Tefferi A, Barbui T. Polycythemia vera: 2024 update on diagnosis, risk-stratification, and management. *Am J Hematol*. 2023;98(9):1465–87.
20. Gisslinger H, Zagrijtschuk O, Buxhofer-Ausch V, Thaler J, Schloegl E, Gastl GA, et al. Ropeginterferon alfa-2b, a novel IFN α -2b, induces high response rates with low toxicity in patients with polycythemia vera. *Blood*. 2015 Oct 8;126(15):1762–9.
21. Mascarenhas J, Kosiorek HE, Prchal JT, Rambaldi A, Berenzon D, Yacoub A, et al. A randomized phase 3 trial of interferon- α vs hydroxyurea in polycythemia vera and essential thrombocythemia. *Blood*. 2022 May 12;139(19):2931–41.
22. Gisslinger H, Klade C, Georgiev P, Krochmalczyk D, Gercheva-Kyuchukova L, Egyed M, et al. Ropeginterferon alfa-2b versus standard therapy for polycythemia vera (PROUD-PV and CONTINUATION-PV): a randomised, non-inferiority, phase 3 trial and its extension study. *Lancet Haematol*. 2020 Mar 1;7(3):e196–208.
23. Kiladjian JJ, Klade C, Georgiev P, Krochmalczyk D, Gercheva-Kyuchukova L, Egyed M, et al. Long-term outcomes of polycythemia vera patients treated with ropeginterferon Alfa-2b. *Leukemia*. 2022 May;36(5):1408–11.
24. Gisslinger H, Klade C, Georgiev P, Krochmalczyk D, Gercheva-Kyuchukova L, Egyed M, et al. Event-free survival in patients with polycythemia vera treated with ropeginterferon alfa-2b versus best available treatment. *Leukemia*. 2023 Oct;37(10):2129–32.
25. Mascarenhas J, Tashi T, El Chaer F, Priego V, Zagrijtschuk O, Qin A, et al. A Phase 3b, Randomized, Open-Label, Parallel Group, Multicenter Study to Assess Efficacy, Safety, and Tolerability of Two Dosing Regimens of Ropeginterferon Alfa-2b-Njft (P1101) in Adult Patients with Polycythemia Vera. *Blood*. 2023 Nov 2;142(Supplement 1):6444.
26. Barosi G, Birgegard G, Finazzi G, Griesshammer M, Harrison C, Hasselbalch H, et al. A unified definition of clinical resistance and intolerance to hydroxycarbamide in polycythemia vera and primary myelofibrosis: results of a European LeukemiaNet (ELN) consensus process. *Br J Haematol*. 2010;148(6):961–3.
27. Barosi G, Besses C, Birgegard G, Briere J, Cervantes F, Finazzi G, et al. A unified definition of clinical resistance/intolerance to hydroxyurea in essential thrombocythemia: results of a consensus process by an international working group. *Leukemia*. 2007 Feb;21(2):277–80.
28. Yacoub A, Mascarenhas J, Kosiorek H, Prchal JT, Berenzon D, Baer MR, et al. Pegylated interferon alfa-2a for polycythemia vera or essential thrombocythemia resistant or intolerant to hydroxyurea. *Blood*. 2019 Oct 31;134(18):1498–509.

29. Vannucchi AM, Kiladjian JJ, Griesshammer M, Masszi T, Durrant S, Passamonti F, et al. Ruxolitinib versus Standard Therapy for the Treatment of Polycythemia Vera. *N Engl J Med*. 2015 Jan 29;372(5):426–35.
30. Passamonti F, Griesshammer M, Palandri F, Egyed M, Benevolo G, Devos T, et al. Ruxolitinib for the treatment of inadequately controlled polycythemia vera without splenomegaly (RESPONSE-2): a randomised, open-label, phase 3b study. *Lancet Oncol*. 2017 Jan 1;18(1):88–99.
31. Harrison CN, Nangalia J, Boucher R, Jackson A, Yap C, O'Sullivan J, et al. Ruxolitinib Versus Best Available Therapy for Polycythemia Vera Intolerant or Resistant to Hydroxycarbamide in a Randomized Trial. *J Clin Oncol*. 2023 Jul;41(19):3534–44.
32. Barbui T, Vannucchi AM, De Stefano V, Carobbio A, Ghirardi A, Carioli G, et al. Ropoginterferon versus Standard Therapy for Low-Risk Patients with Polycythemia Vera. *NEJM Evid*. 2023 May 23;2(6):EVIDoA2200335.
33. Barbui T, Vannucchi AM, Stefano VD, Masciulli A, Carobbio A, Ferrari A, et al. Ropoginterferon alfa-2b versus phlebotomy in low-risk patients with polycythemia vera (Low-PV study): a multicentre, randomised phase 2 trial. *Lancet Haematol*. 2021 Mar 1;8(3):e175–84.
34. Ganz T. Anemia of Inflammation. *N Engl J Med*. 2019 Sep 19;381(12):1148–57.
35. Kremyanskaya M, Kuykendall AT, Pemmaraju N, Ritchie EK, Gotlib J, Gerds A, et al. Rusfertide, a Hepcidin Mimetic, for Control of Erythrocytosis in Polycythemia Vera. *N Engl J Med*. 2024 Feb 21;390(8):723–35.
36. Bankar A, Pettit K, Shatzel J, Yacoub A, Pemmaraju N, Gill H, et al. VERIFY: A randomized controlled phase 3 study of the hepcidin mimetic rusfertide (PTG-300) in patients with polycythemia vera (PV). *J Clin Oncol*. 2024 Jun;42(16_suppl):TPS6592–TPS6592.
37. Barbui T. Appropriate management of Polycythemia Vera with cytoreductive drug therapy: European LeukemiaNet 2021 recommendations. *Hematol Transfus Cell Ther*. 2022 Oct 1;44:S3–4.
38. Barbui T, Barosi G, Birgegard G, Cervantes F, Finazzi G, Griesshammer M, et al. Philadelphia-negative classical myeloproliferative neoplasms: critical concepts and management recommendations from European LeukemiaNet. *J Clin Oncol Off J Am Soc Clin Oncol*. 2011 Feb 20;29(6):761–70.
39. Barbui T, Vannucchi AM, Buxhofer-Ausch V, De Stefano V, Betti S, Rambaldi A, et al. Practice-relevant revision of IPSET-thrombosis based on 1019 patients with WHO-defined essential thrombocythemia. *Blood Cancer J*. 2015 Nov;5(11):e369–e369.
40. Tefferi A, Vannucchi AM, Barbui T. Essential thrombocythemia: 2024 update on diagnosis, risk stratification, and management. *Am J Hematol*. 2024;99(4):697–718.
41. Alvarez-Larrán A, Pereira A, Guglielmelli P, Hernández-Boluda JC, Arellano-Rodrigo E, Ferrer-Marín F, et al. Antiplatelet therapy versus observation in low-risk essential thrombocythemia with a CALR mutation. *Haematologica*. 2016 Aug 1;101(8):926–31.
42. Alvarez-Larrán A, Cervantes F, Pereira A, Arellano-Rodrigo E, Pérez-Andreu V, Hernández-Boluda JC, et al. Observation versus antiplatelet therapy as primary prophylaxis for thrombosis in low-risk essential thrombocythemia. *Blood*. 2010 Aug 26;116(8):1205–10; quiz 1387.
43. Harrison CN, Campbell PJ, Buck G, Wheatley K, East CL, Bareford D, et al. Hydroxyurea Compared with Anagrelide in High-Risk Essential Thrombocythemia. *N Engl J Med*. 2005 Jul 7;353(1):33–45.
44. Harrison CN, Mead AJ, Panchal A, Fox S, Yap C, Gbandi E, et al. Ruxolitinib vs best available therapy for ET intolerant or resistant to hydroxycarbamide. *Blood*. 2017 Oct 26;130(17):1889–97.
45. Verstovsek S, Komatsu N, Gill H, Jin J, Lee SE, Hou HA, et al. SURPASS-ET: phase III study of ropoginterferon alfa-2b versus anagrelide as second-line therapy in essential thrombocythemia. *Future Oncol Lond Engl*. 2022 Sep;18(27):2999–3009.
46. Barbui T, Vannucchi AM, Buxhofer-Ausch V, De Stefano V, Betti S, Rambaldi A, et al. Practice-relevant revision of IPSET-thrombosis based on 1019 patients with WHO-defined essential thrombocythemia. *Blood Cancer J*. 2015;5(11):e369.
47. Gangat N, Tefferi A. Myelofibrosis biology and contemporary management. *Br J Haematol*. 2020;191(2):152–70.
48. Grinfeld J, Nangalia J, Baxter EJ, Wedge DC, Angelopoulos N, Cantrill R, et al. Classification and Personalized Prognosis in Myeloproliferative Neoplasms. *N Engl J Med*. 2018 Oct 11;379(15):1416–30.
49. Cervantes F, Dupriez B, Pereira A, Passamonti F, Reilly JT, Morra E, et al. New prognostic scoring system for primary myelofibrosis based on a study of the International Working Group for Myelofibrosis Research and Treatment. *Blood*. 2009 Mar 26;113(13):2895–901.
50. Passamonti F, Cervantes F, Vannucchi AM, Morra E, Rumi E, Pereira A, et al. A dynamic prognostic model to predict survival in primary myelofibrosis: a study by the IWG-MRT (International Working Group for Myeloproliferative Neoplasms Research and Treatment). *Blood*. 2010 Mar 4;115(9):1703–8.
51. Gangat N, Caramazza D, Vaidya R, George G, Begna K, Schwager S, et al. DIPSS plus: a refined Dynamic International Prognostic Scoring System for primary myelofibrosis that incorporates prognostic information from karyotype, platelet count, and transfusion status. *J Clin Oncol Off J Am Soc Clin Oncol*. 2011 Feb 1;29(4):392–7.
52. Vannucchi AM, Lasho TL, Guglielmelli P, Biamonte F, Pardanani A, Pereira A, et al. Mutations and prognosis in primary myelofibrosis. *Leukemia*. 2013 Sep;27(9):1861–9.
53. Tefferi A, Guglielmelli P, Lasho TL, Gangat N, Ketterling RP, Pardanani A, et al. MIPSS70+ Version 2.0: Mutation and Karyotype-Enhanced International Prognostic Scoring System for Primary Myelofibrosis. *J Clin Oncol*. 2018 Jun 10;36(17):1769–70.
54. Guglielmelli P, Lasho TL, Rotunno G, Mudireddy M, Mannarelli C, Nicolosi M, et al. MIPSS70: Mutation-Enhanced International Prognostic Score System for Transplantation-Age Patients With Primary Myelofibrosis. *J Clin Oncol*. 2018 Feb;36(4):310–8.

55. Passamonti F, Giorgino T, Mora B, Guglielmelli P, Rumi E, Maffioli M, et al. A clinical-molecular prognostic model to predict survival in patients with post polycythemia vera and post essential thrombocythemia myelofibrosis. *Leukemia*. 2017 Dec;31(12):2726–31.
56. Kröger N, Bacigalupo A, Barbui T, Ditschkowski M, Gagelmann N, Grieshammer M, et al. Indication and management of allogeneic haematopoietic stem-cell transplantation in myelofibrosis: updated recommendations by the EBMT/ELN International Working Group. *Lancet Haematol*. 2024 Jan;11(1):e62–74.
57. Maze D, Arcasoy MO, Henrie R, Cerquozzi S, Kamble R, Al-Hadidi S, et al. Upfront allogeneic transplantation versus JAK inhibitor therapy for patients with myelofibrosis: a North American collaborative study. *Bone Marrow Transplant*. 2024 Feb;59(2):196–202.
58. Davidson MB, Gupta V. Application of Stem Cell Therapy in Myelofibrosis. *Hematol Oncol Clin North Am*. 2021 Apr;35(2):391–407.
59. Rajendra A, Gupta V. Advances in Stem Cell Transplantation for Myelofibrosis. *Curr Hematol Malig Rep*. 2024 Sep 6;
60. Harrison C, Kiladjan JJ, Al-Ali HK, Gisslinger H, Waltzman R, Stalbovskaya V, et al. JAK Inhibition with Ruxolitinib versus Best Available Therapy for Myelofibrosis. *N Engl J Med*. 2012 Mar 1;366(9):787–98.
61. Verstovsek S, Mesa RA, Gotlib J, Levy RS, Gupta V, DiPersio JF, et al. A Double-Blind, Placebo-Controlled Trial of Ruxolitinib for Myelofibrosis. *N Engl J Med*. 2012 Mar 1;366(9):799–807.
62. Gupta V, Cerquozzi S, Foltz L, Hillis C, Devlin R, Elsayy M, et al. Patterns of Ruxolitinib Therapy Failure and Its Management in Myelofibrosis: Perspectives of the Canadian Myeloproliferative Neoplasm Group. *JCO Oncol Pract*. 2020 Jul;16(7):351–9.
63. Pardanani A, Harrison C, Cortes JE, Cervantes F, Mesa RA, Milligan D, et al. Safety and Efficacy of Fedratinib in Patients With Primary or Secondary Myelofibrosis: A Randomized Clinical Trial. *JAMA Oncol*. 2015 Aug 1;1(5):643–51.
64. Pardanani A, Tefferi A, Masszi T, Mishchenko E, Drummond M, Jourdan E, et al. Updated results of the placebo-controlled, phase III JAKARTA trial of fedratinib in patients with intermediate-2 or high-risk myelofibrosis. *Br J Haematol*. 2021 Oct;195(2):244–8.
65. Harrison CN, Schaap N, Vannucchi AM, Kiladjan JJ, Tiu RV, Zachee P, et al. Janus kinase-2 inhibitor fedratinib in patients with myelofibrosis previously treated with ruxolitinib (JAKARTA-2): a single-arm, open-label, non-randomised, phase 2, multicentre study. *Lancet Haematol*. 2017 Jul 1;4(7):e317–24.
66. Gupta V, Yacoub A, Verstovsek S, Mesa R, Harrison C, Vannucchi AM, et al. Safety and Efficacy of Fedratinib in Patients with Primary (P), Post-Polycythemia Vera (Post-PV), and Post-Essential Thrombocythemia (Post-ET) Myelofibrosis (MF) Previously Treated with Ruxolitinib: Primary Analysis of the FREEDOM Trial. *Blood*. 2022 Nov 15;140(Supplement 1):3935–7.
67. PRODUCT MONOGRAPH FEDRATINIB HEALTH CANADA [Internet]. [cited 2024 Oct 29]. Available from: https://pdf.hres.ca/dpd_pm/00076169.PDF
68. Mesa RA, Kiladjan JJ, Catalano JV, Devos T, Egyed M, Hellmann A, et al. SIMPLIFY-1: A Phase III Randomized Trial of Momelotinib Versus Ruxolitinib in Janus Kinase Inhibitor-Naïve Patients With Myelofibrosis. *J Clin Oncol*. 2017 Dec;35(34):3844–50.
69. Mesa R, Oh ST, Gerds AT, Gupta V, Catalano J, Cervantes F, et al. Momelotinib reduces transfusion requirements in patients with myelofibrosis. *Leuk Lymphoma*. 2022 Jun 7;63(7):1718–22.
70. Mascarenhas J, Hoffman R, Talpaz M, Gerds AT, Stein B, Gupta V, et al. Pacritinib vs Best Available Therapy, Including Ruxolitinib, in Patients With Myelofibrosis: A Randomized Clinical Trial. *JAMA Oncol*. 2018 May 1;4(5):652–9.
71. Rampal RK, Grosicki S, Chraniuk D, Abruzzese E, Bose P, Gerds AT, et al. Pelabresib in Combination with Ruxolitinib for Janus Kinase Inhibitor Treatment-Naïve Patients with Myelofibrosis: Results of the MANIFEST-2 Randomized, Double-Blind, Phase 3 Study. *Blood*. 2023 Nov 2;142(Supplement 1):628.
72. Pemmaraju N, Mead AJ, Somerville TC, McCloskey JK, Palandri F, Koschmieder S, et al. Transform-1: A Randomized, Double-Blind, Placebo-Controlled, Multicenter, International Phase 3 Study of Navitoclax in Combination with Ruxolitinib Versus Ruxolitinib Plus Placebo in Patients with Untreated Myelofibrosis. *Blood*. 2023 Nov 2;142(Supplement 1):620.
73. Mascarenhas J, Kremyanskaya M, Patriarca A, Palandri F, Devos T, Passamonti F, et al. MANIFEST: Pelabresib in Combination With Ruxolitinib for Janus Kinase Inhibitor Treatment-Naïve Myelofibrosis. *J Clin Oncol*. 2023 Nov 10;41(32):4993–5004.
74. Elena C, Passamonti F, Rumi E, Malcovati L, Arcaini L, Boveri E, et al. Red blood cell transfusion-dependency implies a poor survival in primary myelofibrosis irrespective of IPSS and DIPSS. *Haematologica*. 2011 Jan;96(1):167–70.
75. Gupta V, Harrison C, Hexner EO, Al-Ali HK, Foltz L, Montgomery M, et al. The impact of anemia on overall survival in patients with myelofibrosis treated with ruxolitinib in the COMFORT studies. *Haematologica*. 2016 Dec 1;101(12):e482–4.
76. Cervantes F, Alvarez-Larrán A, Hernández-Boluda JC, Sureda A, Granell M, Vallansot R, et al. Darbepoetin-alpha for the anaemia of myelofibrosis with myeloid metaplasia. *Br J Haematol*. 2006 Jul;134(2):184–6.
77. Cervantes F, Alvarez-Larrán A, Hernández-Boluda JC, Sureda A, Torrealbadell M, Montserrat E. Erythropoietin treatment of the anaemia of myelofibrosis with myeloid metaplasia: results in 20 patients and review of the literature. *Br J Haematol*. 2004 Nov;127(4):399–403.
78. Tsiara SN, Chaidos A, Bourantas LK, Kapsali HD, Bourantas KL. Recombinant human erythropoietin for the treatment of anaemia in patients with chronic idiopathic myelofibrosis. *Acta Haematol*. 2007;117(3):156–61.
79. Passamonti F, Harrison CN, Mesa RA, Kiladjan JJ, Vannucchi AM, Verstovsek S. Anemia in myelofibrosis: Current and emerging treatment options. *Crit Rev Oncol Hematol*. 2022 Dec 1;180:103862.
80. Gerds AT, Harrison C, Kiladjan JJ, Mesa R, Vannucchi AM, Komrokji R, et al. Safety and efficacy of luspatercept for the treatment of anemia in patients with myelofibrosis. *Blood Adv*. 2024 Aug